

# Isolation Categories

Iso can only be changed/discontinued on order of a clinician

Category	Definition	Housing	Movement	Discontinuation of Isolation
Suspect or Positive Case Isolation	Symptoms and awaiting test/result or positive	Single, Solid door with port  Ok to house known positive cases together  Positive cases go to Safety Cell in CJ1	No Court, No Visiting In pod video appointments ok Shower alone Phone at cellside ok with covering over the port  Staff wears N95, face shield and gloves. If close or contact > 10 min or transport, also gown	If negative COVID test ->Other Respiratory Isolation  <b>If positive COVID test</b> 1. Liberalize movement with mask at day 10 (same as other resp iso) 2. Release from iso after 14 days. No additional testing necessary as of 7/8/20.
Other Respiratory Isolation	Negative COVID test, continue isolation to prevent spread of other respiratory viruses	Solid door with port or single celled	Court/Visiting with mask  Staff wears surgical or cloth mask	Symptoms improving and 7 days from symptom onset AND 72 hours fever free; shorter case by case. Lowest risk isolation (can moved out of isolation early if there is a space crunch)
Exposure Isolation	Asymptomatic, high-risk exposure to confirmed case or outbreak or Inpatient/PES/ED return (Does not apply to return from a ZSFG visit for specialty appointment)	Solid door with port or single celled  Ok to house in same cell as same (if same exposure eg outbreak)	Court/Visiting with mask Walk alone or in group of same  Staff wears Surgical or cloth mask	Exposed to confirmed case: 14 days (no shorter) with negative tests at days 7 and 14  Hosp return: 3 days, no additional testing

For all above: phones, showers, tables and any other shared item should be disinfected after use. Patients must use masks when out of bunk/cell and when using phone.

7/9/2020